

Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. In particular, solutions found from a graphic display calculator should be supported by suitable working, e.g. if graphs are used to find a solution, you should sketch these as part of your answer. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. All students should therefore be advised to show their working

1. [Maximum mark: 16]

The function  $f$  is defined by  $f: x \mapsto -0.5x^2 + 2x + 2.5$ .  $f(0) = 5(0) + 2(0) + 2.5 = y$

(a) Write down

(i)  $f'(x)$ ;  $-x + 2$

(ii)  $f'(0)$ ;  $2$

(b) Let  $N$  be the normal to the curve at the point where the graph intercepts the  $y$ -axis. Show that the equation of  $N$  may be written as  $y = -0.5x + 2.5$ .

Let  $g: x \mapsto -0.5x + 2.5$ .

$0 = -x + 2$   $x = 2$   $[2 \text{ marks}]$   
 $50 \perp m = \frac{2}{1}$   $[3 \text{ marks}]$   
 $y = -0.5x + 2.5$

(c) (i) Find the solutions of  $f(x) = g(x)$ .

(ii) Hence find the coordinates of the other point of intersection of the normal and the curve.

$(-5, -20)$

[6 marks]

(d) Let  $R$  be the region enclosed between the curve and  $N$ .

(i) Write down an expression for the area of  $R$ .

(ii) Hence write down the area of  $R$ .

can use calc here

$\int_{-5}^0 (-0.5x^2 + 2x + 2.5) dx - \int_{-5}^0 -0.5x + 2.5 dx$   $[5 \text{ marks}]$

$0.5x^2 + 2x + 2.5 = -0.5x + 2.5$

$0.5x^2 + 2.5x = 0$

$0.5x(x + 5) = 0$

$x = 0$  or  $x = -5$

$\int_{-5}^0 -0.5x^2 + 2.5x dx$   
 $-\left[\frac{1}{6}x^3 + 1.25x^2\right]_{-5}^0$   
 $0 - \left(-\frac{1}{6}(-5)^3 + 1.25(-5)^2\right)$

$+20.83 + 31.25$

$52.08$

$f(-5) = 0.5(-5)^2 + 2(-5) + 2.5$   
 $-12.5 - 10 + 2.5 = -20$

2. [Maximum mark: 18]

A farmer owns a triangular field  $ABC$ . One side of the triangle,  $[AC]$ , is 104 m, a second side,  $[AB]$ , is 65 m and the angle between these two sides is  $60^\circ$ .

(a) Use the cosine rule to calculate the length of the third side of the field. [3 marks]


$$CB^2 = 104^2 + 65^2 - 2(104)(65)\cos 60 = 91$$

(b) Given that  $\sin 60^\circ = \frac{\sqrt{3}}{2}$ , find the area of the field in the form  $p\sqrt{3}$

where  $p$  is an integer.

$$\frac{1}{2}(104)(65)\left(\frac{\sqrt{3}}{2}\right) = 1690\sqrt{3} \quad [3 \text{ marks}]$$

Let  $D$  be a point on  $[BC]$  such that  $[AD]$  bisects the  $60^\circ$  angle. The farmer divides the field into two parts  $A_1$  and  $A_2$  by constructing a straight fence  $[AD]$  of length  $x$  metres, as shown on the diagram below.



$$\sin 30 = \frac{1}{2}$$

(c) (i) Show that the area of  $A_1$  is given by  $\frac{65x}{4}$ .

$$\frac{1}{2}ab\sin C = \frac{1}{2}(x)(65)\left(\frac{1}{2}\right) = \frac{65x}{4}$$

(ii) Find a similar expression for the area of  $A_2$ .

$$\frac{104x}{4}$$

(iii) Hence, find the value of  $x$  in the form  $q\sqrt{3}$ , where  $q$  is an integer. [7 marks]

(d) (i) Explain why  $\sin \hat{ADC} = \sin \hat{ADB}$ . Because they are supplementary

(ii) Use the result of part (i) and the sine rule to show that

$$\frac{BD}{DC} = \frac{5}{8}$$

$$\frac{\sin x}{65} = \frac{\sin(180-x)}{104}$$

$$\sin x = \frac{65}{104} \quad [5 \text{ marks}]$$

$$\frac{104x}{4} + \frac{65x}{4} = 1690\sqrt{3}$$


$$169x = 6760\sqrt{3} =$$

$$x = 40\sqrt{3}$$

Turn over

3. [Total mark: 22]

Part A [Maximum mark: 14]

The diagram below shows the graphs of  $f(x) = 1 + e^{2x}$ ,  $g(x) = 10x + 2$ ,  $0 \leq x \leq 1.5$ 

$$5 = 1 + e^{2a}$$

$$4 = e^{2a}$$

$$\frac{\ln 4}{2} = a$$


- (a) (i) Write down an expression for the vertical distance  $p$  between the graphs of  $f$  and  $g$ .

$$g(x) - f(x) = p$$

- (ii) Given that  $p$  has a maximum value for  $0 \leq x \leq 1.5$ , find the value of  $x$  at which this occurs.

$$\ln \frac{5}{2}$$

The graph of  $y = f(x)$  only is shown in the diagram below. When  $x = a$ ,  $y = 5$ .



$$10x + 2 - 1 - e^{2x} = 10x - e^{2x} + 1$$

[6 marks] find deriv.

$$10 - 2e^{2x} = 0$$

$$-2e^{2x} = -10$$

$$\ln e^{2x} = \ln 5$$

$$2x = \ln 5$$

$$x = \frac{\ln 5}{2}$$

- (b) (i) Find  $f^{-1}(x)$ .

- (ii) Hence show that  $a = \ln 2$ .

$$50 \frac{\ln 4}{2} = a \quad \frac{1}{2} \ln 4 = \ln 2$$

[5 marks]

- (c) The region shaded in the diagram is rotated through  $360^\circ$  about the x-axis. Write down an expression for the volume obtained.


[3 marks]

$$\pi \int_0^{\ln 2} (1 + e^{2x})^2 dx$$

**Part B** [Maximum mark: 8]

Consider the function  $h: x \mapsto \frac{x-2}{(x-1)^2}, x \neq 1$ .

A sketch of part of the graph of  $h$  is given below.



The line (AB) is a vertical asymptote. The point P is a point of inflexion.

- (a) Write down the equation of the vertical asymptote.

$x = 1$

- (b) Find  $h'(x)$ , writing your answer in the form

$$\frac{-x^2 + 4x - 3}{(x-1)^4}$$

$$\frac{a-x}{(x-1)^n}$$

$$\frac{u'v - uv'}{u^2} \Rightarrow \frac{1(x-1)^2 - (x-2)(2(x-1))}{(x-1)^4} = \frac{-2x^2 + 6x - 4}{(x-1)^4}$$

where  $a$  and  $n$  are constants to be determined.

[4 marks]

- (c) Given that  $h''(x) = \frac{2x-8}{(x-1)^4}$ , calculate the coordinates of P.

$(4, 2/9)$

[3 marks]

$$\frac{-(x^2 - 4x + 3)}{(x-1)^4} = \frac{-(x-3)(x-1)}{(x-1)^4}$$

$$2x - 8 = 0$$

$$2x = 8$$

$$x = 4$$

$$h(4) = \frac{4-2}{(4-1)^2} = \frac{2}{9}$$

4. [Maximum mark: 19]

Bag A contains 2 red balls and 3 green balls. Two balls are chosen at random from the bag without replacement. Let  $X$  denote the number of red balls chosen. The following table shows the probability distribution for  $X$ .

$X$	0	1	2
$P(X = x)$	$\frac{3}{10}$	$\frac{6}{10}$	$\frac{1}{10}$

(a) Calculate  $E(X)$ , the mean number of red balls chosen.

$0 \cdot \frac{3}{10} + 1 \cdot \frac{6}{10} + 2 \cdot \frac{1}{10} = \frac{6}{10} + \frac{2}{10} = \frac{8}{10}$  [3 marks]

Bag B contains 4 red balls and 2 green balls. Two balls are chosen at random from bag B.

(b) (i) Draw a tree diagram to represent the above information, including the probability of each event.

(ii) Hence find the probability distribution for  $Y$ , where  $Y$  is the number of red balls chosen.

$Y$	0	1	2
$P(Y)$	$\frac{1}{15}$	$\frac{8}{15}$	$\frac{2}{15}$

A standard die with six faces is rolled. If a 1 or 6 is obtained, two balls are chosen from bag A, otherwise two balls are chosen from bag B.


(c) Calculate the probability that two red balls are chosen.

$\frac{9}{30}$

[5 marks]

(d) Given that two red balls are obtained, find the conditional probability that a 1 or 6 was rolled on the die.

[3 marks]



$$P(A|2R) = \frac{P(A \cap 2R)}{P(2R)}$$

$$= \frac{\frac{1}{30}}{\frac{9}{30}} = \frac{1}{9}$$

## 5. [Maximum mark: 15]

In this question, distance is in kilometers, time is in hours.

A balloon is moving at a constant height with a speed of  $18 \text{ km h}^{-1}$ , in the

direction of the vector  $\begin{pmatrix} 3 \\ 4 \\ 0 \end{pmatrix}$ .  $= \sqrt{9+16} = 5$   $5x = 18/5$   $x = 3.6$

At time  $t=0$ , the balloon is at point B with coordinates  $(0, 0, 5)$ .

(a) Show that the position vector  $b$  of the balloon at time  $t$  is given by

$$b = \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 5 \end{pmatrix} + t \begin{pmatrix} 10.8 \\ 14.4 \\ 0 \end{pmatrix} \quad b = \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 5 \end{pmatrix} + t \begin{pmatrix} 3.6 \\ 4.8 \\ 0 \end{pmatrix} \begin{pmatrix} 3 \\ 4 \\ 0 \end{pmatrix} \text{ marks}$$

At time  $t=0$ , a helicopter goes to deliver a message to the balloon. The position vector  $h$  of the helicopter at time  $t$  is given by

$$h = \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 49 \\ 32 \\ 0 \end{pmatrix} + t \begin{pmatrix} -48 \\ -24 \\ 6 \end{pmatrix}$$

(b) (i) Write down the coordinates of the starting position of the helicopter.  $(49, 32, 0)$

(ii) Find the speed of the helicopter.  $\sqrt{(-48)^2 + (-24)^2 + 6^2} = 54 \text{ km h}^{-1}$  [4 marks]

(c) The helicopter reaches the balloon at point R.

(i) Find the time the helicopter takes to reach the balloon.  $5/6 \text{ hr.}$

(ii) Find the coordinates of R.  $(9, 12, 5)$  [5 marks]

$$\begin{aligned} 49 - 48t &= 0 + 10.8 \\ 32 - 24t &= 0 + 14.4 \\ 0 + 6t &= 5 + 0 \end{aligned}$$

$$\begin{aligned} 49 - 48(5/6) &= 9 \\ 32 - 24(5/6) &= 12 \\ 0 + 6(5/6) &= 5 \end{aligned}$$

$$t = 5/6$$